

Timing of flowering and pod initiation in chickpeas

■ By Neroli Graham¹, Rosy Raman², Annie Warren¹ and Muhuddin Anwar²

CHICKPEAS are suited to environmental conditions of many grain growing regions of Australia and are an important winter pulse crop for farming systems. But those same environmental conditions, and other factors, often conspire to significantly reduce yield. For example in northern NSW and southern Queensland, abiotic and biotic factors are estimated to cause yield reductions of 1.7–2.7 tonnes per hectare (Yield Gap Australia, 2018). In 2016, cool spring weather was estimated to have resulted in yield losses of 0.5–0.7 tonnes per hectare in northwest NSW.

Under ideal environmental conditions, chickpea will produce pods within a couple of days of flowering. But if temperatures stay below optimal levels the length of time between the beginning of flowering and the commencement of podding can be more than two months. Furthermore, if average daily temperatures are below 15°C during flowering, there is reduced pollen viability, delayed stigma development and flower drop. Trials and other observations have shown that average temperatures during flowering significantly influenced flower abortion.

Siddique and Sedgley (1986) found that when chickpeas were sown early and experienced average temperatures of 12.5°C during flowering there was 800 aborted flowers per m²,

compared to 0 aborted flowers per m² from a later sowing date where the average temperature was 16.8°C during flowering.

How did we get here?

In 2020, detailed phenology experiments were conducted at Tamworth in northern NSW and Wagga Wagga in southern NSW. In this series of experiments, sowing time was used to assess the capacity of a range of genotypes to produce pods and seeds under cool temperatures. These experiments included a set of 12 diverse genotypes, both released cultivars and advanced breeding lines, and targeted three sowing windows, with an early (late April to early May), main (mid-May to early June) and late season (mid to late June) sowing.

Plant phasic development stages, between late vegetative and pod initiation were defined and dates were recorded.

Changes in phase development that are important for pod and seed development under suboptimal environmental conditions included:

- The date when plants become reproductive;
- The date when 50 per cent of plants have one open flower; and,
- The date of pod initiation.

CROPLANDS

LIGHT YEARS AHEAD


DETECT **TARGET** **ERADICATE** **SAVE**

When you invest in a WEED-IT from Croplands you are also investing in our spot spraying expertise. Whether it's technology or application, our team has the experience to provide you the unparalleled service and ongoing technical support that you need.

Get outstanding results, even on hard-to-kill weeds, with WEED-IT – the world's number one spot sprayer.

WEED-IT precision spraying

FIGURE 1: Phase development of four commercially released cultivars and an early flowering genotype at Tamworth in 2020 for three sowing dates



Vertical line represents the date at which average air temperature rose above 15°C on September 3.

What did we find?

Flowering and pod initiation

Sowing date and location were found to influence the timing of phasic development of individual genotypes. At Tamworth, from the early sowing, in late April (Figure 1), the early flower CBA line had at least one flower on 50 per cent of plants by July 28, 22 days earlier than the next cultivar, CBA Captain and 35

days earlier than the slowest variety Kyabra.

When sown in late April, Kyabra flowered on September 1, two days before the average air temperature rose above 15°C on September 3 (Figure 1). In contrast, when the cultivars were sown in the optimal/main sowing window in late May, the early flower CBA line commenced flowering on August 29, with CBA Captain and PBA Striker achieving 50 per cent flowering on September 5, and Kyabra and PBA Seamer flowering on September 8 (Figure 1).

In contrast to the Tamworth findings, flowering did not commence at Wagga Wagga until September 6, the day the average air temperature rose above 15°C for the first time (Figure 2). The first line to flower from the main sowing in mid-May was the early flower CBA line. Time to flowering was condensed over the

three sowing dates (Figure 2). Date of pod initiation at Wagga Wagga was also condensed, with a small but significant delay in pod initiation for Kyabra when compared to the early flower CBA line at each of the sowings (Figure 2).

Pod initiation, averaged over all genotypes at Tamworth for late April, late May and late June sowing dates, respectively,



Get Ready for **MORE!**

More Capacity. More Speed. More Flex. The all-new FD2 gives you more of everything you need from the company that brought you the Original FlexDraper®.

*Compared to previous model.



20%
More Capacity*



30%
More Speed*



70%
More Flex*




FD2 FlexDraper®

SEE IT IN ACTION!

Find your local dealer at MacDon.com



FIGURE 2: Phase development of four commercially released cultivars and an early flowering genotype at Wagga Wagga in 2020 at three sowing dates



Vertical line represents the date at which average air temperature rose above 15°C on September 6.

occurred 10, 13 and 32 days after average air temperature became optimal (September 3) (Figure 1). The delay in pod initiation, for the late June sowing at Tamworth, reflected the dry spring and depleted soil moisture in dryland cropping systems in northern NSW in 2020.

In contrast, when averaged over all genotypes, at Wagga Wagga, pod initiation for the late April, mid-May and early June sowings, respectively occurred 16, 18 and 25 days, after optimal average air temperature was achieved on September 6, (Figure 2). Pod initiation at Wagga Wagga was in part delayed due to the higher level of temperature fluctuation in September, when temperatures rose and fell, above and below the optimal average air temperature of 15°C.

Relationship between flowering and podding

The relationships between date of 50 per cent flowering and date of pod initiation varied between the two sites. Genotypes sown early at Tamworth, had a spread of dates for the time to 50 per cent flowering between July 19 to September 9 (52 days), while pod initiation occurred quickly, from September 9 to September 13 (4 days).

In comparison, the second and third sowing dates both had a variation in the time to 50 per cent flowering of 17 days and 20 days, while pod initiation was spread over 9 days and 12 days, respectively.

The relationships between date to 50 per cent flowering and date of pod initiation, were compressed at Wagga Wagga when compared to Tamworth. There was overlap of both the date to 50 per cent flower and the date of pod initiation between the three sowing times. Early and main sowing had similar dates to 50 per cent flowering from September 7 to September 18 and pod initiation from September 14 to October 6. It is surmised, that this compression of dates for these phenological stages, may have been due to the greater variation in spring temperatures observed at Wagga Wagga, when compared to Tamworth.

Indicative yield response

When pooled across all genotypes, highest grain yield was observed when crops were sown between mid to late May, yielding 2.5 tonnes per hectare at Tamworth and 2.7 tonnes per hectare at Wagga Wagga. Sowing earlier than mid-May caused a yield loss of approximately 7 per cent at Tamworth and 18 per

cent at Wagga Wagga. In part, the lower grain yield associated with early sowing, may be due to the longer time spent during the reproductive period in suboptimal temperatures.

Individual varieties showed different patterns of yield rankings over the three sowing dates at both Wagga Wagga and Tamworth. Differences in yields between the three sowing dates for four genotypes, CBA Captain, early flower CBA line, Kyabra and PBA Seamer, at Wagga Wagga were significant, whilst at Tamworth they were not. When each of the four genotypes were sown early at Wagga Wagga there was a reduction in yield of between 12 per cent to 14 per cent, when compared to their main season sowing yields.

This loss in grain yield when crops were sown earlier than optimal can

be partially explained by plants developing flowers that do not form pods or pods with seeds when temperatures are suboptimal. Current research, within this project, aims to look at the level of lost flowering opportunities and potential pods and seeds, when crops flower under suboptimal conditions, through measuring the conversion of reproductive nodes into pods and pods that contain seeds.

To sum up

Matching developmental phases – in particular flowering and pod initiation – to optimal environmental conditions increases the yield potential of chickpea cultivars. Importantly, results from these experiments show that early flowering does not translate to earlier pod initiation.

This was particularly highlighted when comparing genotypes from an early sowing date at Tamworth in 2020.

But the results from these experiments do indicate that there is greater variation in pod initiation between genotypes from main and late season sowings, with earlier flowering genotypes initiating pods earlier than later flowering lines.

It can be seen from these experiments that early flowering does not equal early pod initiation or increased grain yield.

Authors: 1 NSW DPI, Tamworth and 2 NSW DPI, Wagga Wagga.

The research undertaken as part of this project is made possible by the significant contributions of growers through both trial cooperation and the support of the GRDC, the author would like to thank them for their continued support.

The author would like to thank current and past staff that have worked on this project: Annie Warren and Helene Davidson at Tamworth and Rosy Raman, Muhuddin Anwar, Mark Richards, Jess Simpson, Asad Asaduzzaman and Joe Moore at Wagga Wagga.

The author would like to acknowledge the assistance of other research teams at both TAI and WWAI for trial management these include CBA lead Kirsty Hobson at TAI, including Judy Duncan, Andrew George, Ben Frazer, Belinda Rowe, Mandy Rowe and Madonna Rowe, winter cereal team lead at TAI by Rick Graham including Stephen Morphett, Peter Formann Rod Bambach, Bailey Skewes and Mick Dal Santo and pulse evaluation team at WWAI lead by Mark Richards including Michael McCaig, Ollie Owens and Karl Moore.

The author would also like to thank the farm managers: Scott Goodwin at Liverpool Plains Research Station, Breeza, Ross Kamphorst at Glen Innes Advisory and Research Station, and the farm managers and staff at Tamworth Agricultural Institute and Wagga Wagga Agricultural Institute.

Contact details: Dr Neroli Graham NSW Department of Primary Industries 4 Marsden Park Road, Calala, NSW, 2340 Ph: 02 6763 1274 Email: neroli.graham@dpi.nsw.gov.au